



P04-15. Prevalence of Broadly Neutralizing Antibody Responses During Acute/Early HIV Infection

Citation

Mikell, I., D. Sather, S. Kalams, Marcus Altfeld, and L. Stamatatos. 2009. P04-15. Prevalence of broadly neutralizing antibody responses during acute/early HIV infection. *Retrovirology* 6(Suppl 3): P43.

Published Version

doi:10.1186/1742-4690-6-S3-P43

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:4774089>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Poster presentation

P04-I5. Prevalence of broadly neutralizing antibody responses during acute/early HIV infection

I Mikell^{*1}, D Sather¹, S Kalams², M Altfeld³ and L Stamatatos¹

Address: ¹Seattle Biomedical Research Institute, Seattle, WA, USA, ²Department of Microbiology and Immunology, Vanderbilt University School of Medicine, Nashville, TN, USA and ³Partners AIDS Research Center, Massachusetts General Hospital, Harvard, Boston, USA

* Corresponding author

from AIDS Vaccine 2009
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, **6**(Suppl 3):P43 doi:10.1186/1742-4690-6-S3-P43

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P43>

© 2009 Mikell et al; licensee BioMed Central Ltd.

Background

Determining how cross-reactive neutralizing antibody (NAb) responses develop during natural HIV-1 infection may provide key information to understand the role they play in controlling the infection and in disease progression. Here we investigated the frequency and breadth of broadly NAb responses during acute and early infection in a well controlled cohort, and attempted to characterize the factors associated with the development of such responses.

Methods

The plasma of 38 clade B acutely-infected, antiretroviral-naïve subjects from two cohorts was screened for breadth of neutralization against a panel of heterologous isolates in an Env pseudovirus neutralization assay. Clade A, B and C variants were chosen from reference panels of viruses, created to evaluate the NAb responses elicited during infection or immunization.

Results

Our preliminary screening strategy demonstrated that broadly neutralizing antibodies can be detected in a third of the infected subjects. Breadth of neutralization can develop as early as one year during natural HIV infection, however the majority of breadth was observed at over two years post infection.

Conclusion

Cross-reactive NAb responses are developed more frequently during early infection than previously thought. The consequences

of this 'early' development of cross-reactive NAb responses on plasma viremia and disease progression are under investigation.

The work described was supported by National Institute of Health grant number R01 AI047708-11.